

Scoping Review of Racial, Ethnic, and Sex Disparities in the Diagnosis and Management of Hemorrhagic Stroke

Lakshman Kalasapudi, BA, MA, Stacey Williamson, MD, Andrea G. Shipper, MSLIS, Melissa Motta, MD, Charles Esenwa, MD, Fadar Oliver Otite, MD, Seemant Chaturvedi, MD, and Nicholas A. Morris, MD

Neurology® 2023;101:e267-e276. doi:10.1212/WNL.0000000000207406

Correspondence

Dr. Morris
nicholas.morris@som.umaryland.edu

Abstract

Background and Objectives

In the United States, Black, Hispanic, and Asian Americans experience excessively high incidence rates of hemorrhagic stroke compared with White Americans. Women experience higher rates of subarachnoid hemorrhage than men. Previous reviews detailing racial, ethnic, and sex disparities in stroke have focused on ischemic stroke. We performed a scoping review of disparities in the diagnosis and management of hemorrhagic stroke in the United States to identify areas of disparities, research gaps, and evidence to inform efforts aimed at health equity.

Methods

We included studies published after 2010 that assessed racial and ethnic or sex disparities in the diagnosis or management of patients aged 18 years or older in the United States with a primary diagnosis of spontaneous intracerebral hemorrhage or aneurysmal subarachnoid hemorrhage. We did not include studies assessing disparities in incidence, risks, or mortality and functional outcomes of hemorrhagic stroke.

Results

After reviewing 6,161 abstracts and 441 full texts, 59 studies met our inclusion criteria. Four themes emerged. First, few data address disparities in acute hemorrhagic stroke. Second, racial and ethnic disparities in blood pressure control after intracerebral hemorrhage exist and likely contribute to disparities in recurrence rates. Third, racial and ethnic differences in end-of-life care exist, but further work is required to understand whether these differences represent true disparities in care. Fourth, very few studies specifically address sex disparities in hemorrhagic stroke care.

Discussion

Further efforts are necessary to delineate and correct racial, ethnic, and sex disparities in the diagnosis and management of hemorrhagic stroke.

MORE ONLINE

 **CME Course**
[NPub.org/cmelist](https://www.npub.org/cmelist)

Glossary

AHA = American Heart Association; ASA = American Stroke Association; DNR = do not resuscitate; ED = emergency department; ERICH = Ethnic/Racial Variations of Intracerebral Hemorrhage; EVD = external ventricular drain; GWTG-S = Get With The Guidelines–Stroke; ICH = intracerebral hemorrhage; ICP = intracranial pressure; IVH = intraventricular hemorrhage; NIS = Nationwide Inpatient Study; PEG = percutaneous endoscopic gastrostomy; SAH = subarachnoid hemorrhage.

Attention to racial, ethnic, and sex disparities in the diagnosis and management of acute ischemic stroke is surging.^{1,2} The focus is shifting from documenting disparities in care to seeking solutions and evaluating interventions.³ Less is known about disparities in the diagnosis and management of hemorrhagic stroke, perhaps because ischemic strokes outnumber hemorrhagic strokes in the United States by nearly 7:1.⁴ Hemorrhagic stroke, however, has substantially higher short-term and long-term mortality compared with ischemic stroke⁵ and disproportionately afflicts Black, Hispanic, and Asian Americans compared with White Americans.⁶ Furthermore, the incidence of intracerebral hemorrhage (ICH), the predominant form of hemorrhagic stroke, is increasing.⁷ Meanwhile, aneurysmal subarachnoid hemorrhage (SAH) disproportionately afflicts women. It is therefore important to understand and address disparities in both ischemic and hemorrhagic stroke. To address this gap, we performed a scoping review of racial, ethnic, and sex disparities pertaining to the diagnosis and management of hemorrhagic stroke. We aimed to identify targets for future research with the goal of improving health equity for hemorrhagic stroke patients.

Methods

We developed a search strategy focusing on racial and ethnic or sex disparities regarding the diagnosis or management of hemorrhagic stroke. We chose a scoping review design to map the literature on the topic and identify key gaps in current research. Scoping reviews differ from systematic reviews because they do not attempt to answer a specific question or inform practice; rather, they identify important concepts or knowledge gaps, delineate the available evidence, and examine how research is conducted in a field.⁸ We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews guideline.⁹

Standard Protocol Approvals, Registrations, and Patient Consents

This scoping review was not registered. As a scoping review of published literature, we did not seek institutional review board approval.

Eligibility Criteria

We included English language studies that assessed racial and ethnic or sex disparities as either a primary or secondary outcome in the diagnosis or management of patients aged 18 years or older in the United States with a primary diagnosis of

spontaneous ICH or aneurysmal SAH. We limited our search to studies after 2010 to accurately portray the current state of disparities. We did not include studies assessing disparities in incidence, risks, or mortality and functional outcomes of hemorrhagic stroke. We excluded studies that did not present results by stroke subtype and disparities by race/ethnicity or sex. Of note, we treated all sex and gender differences as assessments of biologic sex differences because these 2 terms were conflated in most studies, and there were no specific references to gender identity.

Search Strategy and Study Selection

On September 1, 2020, and February 22, 2022, we searched EMBASE, PubMed, and Cochrane Register of Controlled Trials (full search strategy in eAppendix 1, links.lww.com/WNL/C824). We also included any relevant articles found in references of included studies that were not captured in the search. Two authors (L.K., N.A.M.) independently reviewed all abstracts that met the search criteria. In case of disagreement, a third author (M.M.) served as arbiter. We repeated this for a full-text review. We used American Heart Association/American Stroke Association (AHA/ASA) guidelines for the management of ICH and SAH as a framework for reporting our findings, but not to limit the search or grade the strength of the evidence.^{10,11} When possible, we conformed to the AHA/ASA guidelines on disparities research.¹²

Data Availability

A full list of studies excluded from the review made by the authors will be available on request.


Results

Our search yielded 6,161 unique studies for screening, and 59 studies met eligibility criteria (Figure; eTable 1, links.lww.com/WNL/C826). Forty-two studies assessed disparities among patients with ICH; 17 studies assessed disparities among patients with SAH. AHA/ASA guideline content areas without any studies meeting study eligibility criteria are summarized in Tables 1 and 2.

Intracerebral Hemorrhage

Of the 42 studies assessing disparities in patients with ICH, 29 studies assessed racial and ethnic disparities, 5 studies assessed sex disparities, and 8 studies assessed both racial and ethnic and sex disparities. Thirty of 42 studies demonstrated significant disparities or differences in the evaluation or medical management of patients with ICH.

Figure PRISMA Flowchart of Study Selection



After reviewing 6,161 study abstracts for eligibility, we assessed 444 full-text studies, 59 of which met eligibility criteria for review. PRISMA = Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

Prehospital and Initial Systems of Care/Diagnostic Assessment

No studies assessed for disparities in prehospital management of patients with ICH. Seven studies assessed emergency department (ED) management care. A retrospective analysis of 543 patients from the National Hospital Ambulatory Medical Care Survey–Emergency Department database that included both ischemic and hemorrhagic stroke types found that Black patients, but not Hispanic patients, had significantly longer ED wait times when they were being evaluated for a possible stroke when compared with White patients.¹³ A case-control analysis of 123,623 patients from the Get With The Guidelines–Stroke (GWTG-S) registry found that Black patients were less likely than White patients to have a door to CT time of ≤ 25 minutes after controlling for patient-level and hospital-level characteristics.⁶ By contrast, a retrospective review of 459 patients with ICH from a single center found that race and ethnicity had no effect on time to bed request or time to admission.¹⁴ Conflicting data exist regarding whether Black patients are more likely or not to be screened for drugs of abuse after ICH compared with White patients.^{15,16} One retrospective study found that male patients with ICH were more likely than female patients with ICH to undergo urine drug screening.¹⁶

A retrospective single-center study of 265 patients found that providers activated stroke codes less for female patients presenting with ICH to the ED compared with that for male patients.¹⁷ Last, a multivariable analysis of 205 patients from the Genetic and Environmental Risk Factors for Hemorrhagic Stroke study found that Black patients were more likely to be transferred to a tertiary center than non-Black patients presenting to non-tertiary EDs.¹⁸ Disparities in advanced neuroimaging aimed at

identifying underlying pathogenesis such as CT angiography/venography, MRI, or catheter angiography have not been investigated.

Medical and Neurointensive Treatment

Acute Blood Pressure Lowering

One study scrutinized disparities in acute blood pressure management. A multivariate analysis of 2,069 patients from the Ethnic/Racial Variations of Intracerebral Hemorrhage (ERICH) study, a prospective multicenter case-control study of ICH among Black patients, Hispanic patients, and White patients found that Hispanic patients and Black patients had higher prehospital and ED blood pressure compared with White patients and that Black patients continued to have higher blood pressure than both Hispanic patients and White patients 24 hours after admission despite more often receiving continuous antihypertensive infusions.¹⁹ We did not find any studies investigating sex disparities in acute blood pressure lowering after ICH.

Anticoagulant-Related Hemorrhage/Antiplatelet-Related Hemorrhage/General Hemostasis

We did not find any studies investigating disparities in the reversal of anticoagulation in patients with anticoagulant-related ICH. In a multiple logistic regression model including more than 2,500 patients with ICH from the ERICH study, Black patients with antiplatelet-related ICH were less likely to receive platelet transfusions after ICH.²⁰ This analysis was performed before the PATCH study, which found no benefit and possible harm from platelet transfusion.²¹ No studies explored sex disparities in hemostasis.

Table 1 Overview of Disparities in the Diagnosis and Management of ICH

Component of care	Evidence of disparity?	Key citation(s)	Main results summary	Unstudied aspects
Prehospital and initial systems of care/diagnostic assessment	Yes	#6, #12, #13–17	Black patients had longer ED wait times and longer times from door to CT. Conflicting data re: whether Black patients screened more often for drugs of abuse than White patients. Stroke codes activated less frequently for women presenting with ICH.	Prehospital management, other laboratory testing, and advanced neuroimaging.
Acute blood pressure lowering	Yes	#18	Black patients continued to have higher blood pressure than both Hispanic patients and White patients 24 h after admission.	Blood pressure variability. Sex Disparities.
Hemostasis	Yes	#19	Black patients with antiplatelet-related ICH received less platelet transfusions.	Reversal of anticoagulation, general use of hemostatic agents. Sex disparities in hemostasis.
General medical management	Yes	#6, #21–26	Black patients received DVT prophylaxis more often. Asian, Hispanic, and Black patients screened for dysphagia screening more than White patients. Black and Hispanic patients received more PEGs than White patients.	Systemic anticoagulation/IVC filter placement with symptomatic DVT or PE, care setting, nursing care, glucose management, temperature management.
Seizures and antiseizure medications	No	#27–28	No difference in the utilization of antiseizure medications when controlling for ICH location.	Utilization of continuous EEG, Treatment of seizures. Sex disparities.
Neuroinvasive monitoring, ICP, edema treatment, surgical interventions	Uncertain	#23, #25, #29–30	Higher rates of EVD utilization in Black patients were due to higher rates of IVH. Conflicting findings for craniotomy/craniectomy. Higher rates of EVD utilization in men compared with women.	Therapies for ICP/edema.
Outcome prediction and goals of care	Yes	#6, #31–43	Most (but not all) studies suggest less care limitations and palliative care utilization in Black patients and Hispanic patients. Conflicting results, but overall support for earlier DNR orders placed in women.	Qualitative explorations into reasons for disparities.
Recovery and rehabilitations	Yes	#6, #46	Black patients more likely than White patients to be discharged to an inpatient rehabilitation facility.	Recovery trajectories, rehabilitation quality/intensity. Sex disparities.
Secondary prevention	Yes	#47–49	Black patients and Hispanic patients had a higher risk of ICH recurrence than White patients and worse long-term blood pressure control.	Resumption of antithrombotic agents, lifestyle modifications, patient/caregiver education. Sex disparities.

Abbreviations: DNR = do not resuscitate; DVT = deep vein thrombosis; ED = emergency department; ICH = intracerebral hemorrhage; ICP = intracranial pressure; IVC = inferior vena cava; PE = pulmonary embolism; PEG = percutaneous endoscopic gastrostomy.

Inpatient Care Setting/General Medical Management

The AHA/ASA ICH guideline covers multiple aspects of inpatient care including care setting, prevention and management of medical complications, thromboprophylaxis and treatment of thrombosis, nursing care, glucose management, and temperature management. No studies evaluated care setting, nursing care, glucose management, or temperature management.

In the GWTG-S registry, Black patients were more likely than White patients to receive deep vein thrombosis prophylaxis.⁶ No studies assessed racial, ethnic, or sex disparities in the use of pneumatic boot use or systemic anticoagulation/inferior vena cava filter placement in patients with ICH and symptomatic deep vein thrombosis or pulmonary embolism. Compared with White patients, Asian, Hispanic, and Black patients were more

likely to receive dysphagia screening to prevent pneumonia in the GWTG-S registry.⁶ Sex differences in dysphagia screening have not been examined. In a prospective cohort study of 512 patients from 1 hospital, more non-Hispanic Black patients received percutaneous endoscopic gastrostomy (PEG) tubes than Hispanic or non-Hispanic White patients, but this study did not control for confounders.²² In the same study, higher socioeconomic status was associated with lower odds of pursuing PEG only among Hispanic White patients, suggesting that economic factors may moderate decision-making among different races or ethnicities.

Three retrospective studies using Nationwide Inpatient Study (NIS) data have investigated racial and ethnic PEG placement disparities. A 2016 study of 49,946 ICH admissions found

Table 2 Overview of Disparities in the Diagnosis and Management of SAH

Component of care	Evidence of disparity?	Key citation(s)	Main results	Unstudied aspects
Rehabilitation and recovery	Yes	#67	Black patients more likely to be discharged to postacute care than White, Hispanic, or Asian patients.	Recovery trajectories, rehabilitation quality/intensity. Sex disparities.
Diagnosis	Yes	#51	Non-Hispanic White patients were more likely to be misdiagnosed with nonspecific headache diagnoses.	Utilization of lumbar puncture, neuroimaging including digital subtraction angiography. Sex disparities.
Medical measures to prevent rebleeding	No	#52	No racial, ethnic, or sex disparities in a small study of hyperacute blood pressure management.	Administration of antifibrinolytic agents.
Surgical and endovascular management of aneurysms	Yes	#53–57	Conflicting results. Some studies suggest that Hispanic patients and Asian patients, but not Black patients had delays in aneurysm securement compared to White patients. Men had more delays in aneurysm securement than women.	Advanced aneurysm management (flow diversion, rates of complete aneurysm obliteration).
Hospital characteristics and systems of care	Yes	#63	Black patients and Hispanic patients transferred less than White patients. No sex disparity in transfer likelihood.	Hospital SAH volume, hospital complication rates, hospital credentialing processes
Management of seizures	—	—	—	No recent studies.
Management of vasospasm/delayed cerebral ischemia	—	—	—	No recent studies.
Anesthetic management	—	—	—	Completely unstudied.
Management of medical complications	—	—	—	Completely unstudied.
Management of hydrocephalus	—	—	—	Completely unstudied.
Withdrawal of life-sustaining therapies, palliative care	Yes	#64–66	Black and Hispanic patients were less likely to have comfort-measures-only orders. Women had care withdrawn more often than men.	Qualitative explorations into reasons for disparities.
Family member screening	—	—	—	Completely unstudied.

Abbreviation: SAH = subarachnoid hemorrhage.

non-White patients were more likely to receive a PEG than White patients, especially in smaller hospitals with lower ICH volumes.²³ That finding was confirmed in a separate multivariable analysis from the NIS.²⁴ Another NIS study of 38,080 ICH admissions corroborated the findings but found that hospitals that had $\geq 50\%$ of patients admitted with ICH of either Black race or Hispanic ethnicity did not have higher rates of PEG placement.²⁵ Regarding sex disparities, an analysis from the ERICH study of 2,964 patient records did not find any differences in the placement of PEG tubes.²⁶ In a study of 36,019 admissions from the NIS, minority race/ethnicity (e.g., Hispanic, Black, Asians/Pacific Islander) and male sex were associated with increased odds of nonbeneficial PEG tube placement, with minority male patients more likely than White female patients to receive PEG tube placement.²⁷

Seizures and Antiseizure Medications

An investigation of 744 patients from the ERICH study did not find any racial, ethnic, or sex disparities in the receipt of

prophylactic antiseizure medications.²⁸ An analysis of 3,422 patients from a multicenter prospective cohort found that White patients were twice as likely as Black patients to receive seizure prophylaxis after ICH, but the association was eliminated in a single-center subset controlling for hematoma location.²⁹

No studies evaluated for disparities in the application of continuous EEG or utilization of antiseizure medications after seizure. Sex disparities in the utilization of antiseizure medications after ICH have not been investigated.

Neuroinvasive Monitoring, ICP, Edema Treatment, and Surgical Interventions

We found 4 studies addressing disparities in the procedural recommendations of the AHA/ASA ICH guideline: intracranial pressure (ICP) monitoring and treatment, intraventricular hemorrhage (IVH) management, and surgical treatment.

The first study from the NIS of more than 700,000 patients found higher unadjusted rates of external ventricular drain

(EVD) utilization and craniectomy among all minority races and ethnicities when compared with White patients. After multivariable adjustment, EVD placement was significantly higher in Black patients than White patients, while craniectomy utilization was higher in Asian or Pacific Islander patients, Hispanic patients, and other patients (but not Black patients) compared with White patients.²⁴ A second retrospective study analyzing 226 Black and White patients from a single institution found that Black patients were more likely to receive an EVD due to higher rates of IVH and hydrocephalus.³⁰ An analysis of 2,947 patients from the ERICH study found that there was no significant difference in the use of surgical intervention such as craniotomy or minimally invasive hematoma evacuation between Black, White, and Hispanic patients.³¹ A second study from ERICH found that EVD utilization, but not other surgical procedures, was increased in male patients with ICH compared with female patients with ICH after adjusting for age, hematoma volume, and hematoma location.²⁶ No studies investigated disparities in ICP monitoring through parenchymal probes or edema therapies.

Outcome Prediction and Goals of Care

More than one-third of the included studies addressed end-of-life management in ICH. Eleven evaluated racial or ethnic differences, 4 evaluated sex differences, and 3 evaluated both. Fifteen of 18 articles demonstrated differences in end-of-life care between racial and ethnic groups or between sexes. Twelve studies found that White patients opted for fewer life-sustaining interventions than Black patients.^{6,32-42} Prevalence of care limitations was also low among Hispanic patients.^{6,33,35-37,39-42}

Seven studies explicitly included Asian patients. Five studies found that Asian patients have a higher use of life-sustaining interventions than White patients. Three of these studies grouped Asians with other non-White patients to demonstrate that minorities are more likely to use life-sustaining interventions compared with White patients.^{33,39,40} Three studies found that Asian patients received less palliative care than White patients.^{6,35,36} A single-center study of 396 patients from Hawaii found that there were no differences in the withdrawal of life-sustaining therapies between non-Hispanic White patients and Native Hawaiian and other Pacific Islander patients after adjusting for age.⁴³

A 2017 study from the NIS of 46,735 admissions found that minority-serving hospitals (e.g., hospitals with >50% minority patients) used less palliative care services regardless of the race of the patient when compared with predominantly White-serving hospitals (e.g., hospitals with <25% minority patients).³⁹ In addition, the same study noted that racial and ethnic minorities received less palliative care than White patients independent of hospital status as minority serving or not.

Studies of sex disparities in end-of-life care have reached disparate conclusions. A single-center prospective study of 245

patients from 2013 found that new do-not-resuscitate (DNR) orders were placed more often for female patients with ICH than for male patients with ICH in the first 24 hours of admission despite no significant differences in clinical severity, hemorrhage volume, or medical comorbidities.⁴⁴ A retrospective single-center study of 372 patients from 2013 confirmed that after adjustment for relevant variables, early DNR orders were more likely for female patients than for male patients.⁴⁵ A national analysis of 176 hospitals in the United States found that among 8,069 patients with ICH, female patients were more likely to be discharged to hospice after controlling for age.⁴⁶ These findings were further supported by 2 large NIS studies that found increased DNR and palliative care use in female patients compared with those in male patients.^{35,41} Smaller, more recent single-center studies have not found sex differences in the use or timing of DNR orders^{36,40} nor did an ERICH study looking at DNR, palliative care, and hospice use after controlling for ICH score, prior stroke, and dementia.²⁶

Post-ICH Recovery, Rehabilitation, and Complications

Results from the GWTG-S registry indicated that Black patients were more likely than White patients to receive a rehabilitation assessment during their hospital stay. Black patients, but not Hispanic or Asian patients, were less likely to be discharged to an inpatient rehabilitation facility compared with White patients.⁶

Secondary Prevention

Three studies found that Black patients are at a higher risk of ICH recurrence compared with White patients.⁴⁷⁻⁴⁹ One of those studies, using records from a single center and the ERICH study found that among 2,291 patients, Hispanic patients are also at an increased risk,⁴⁷ while a longitudinal analysis of prospectively collected claims data from 31,355 patients at nonfederal California hospitals found that Asian patients and not Hispanic patients are at an increased risk.⁴⁸ A single-center prospective cohort study of 1,305 patients found that non-White patients were at a higher risk of uncontrolled hypertension at 3 months when compared with White patients, partially due to inadequately prescribed antihypertensive medication.⁴⁹ A prospective cohort study of 162 patients found that Black patients were more likely to have stage I/II hypertension 1 year after ICH than White patients despite showing no difference in hypertension at 30 days.⁵⁰ Analysis from a combined single-center and ERICH cohort found that Black and Hispanic patients had more hypertension at follow-up, but this finding only partially explained their excess ICH recurrence risk.⁴⁷ No studies investigated sex disparities in blood pressure management after ICH nor disparities in the management of medications including antithrombotic agents, lifestyle modifications, or patient/caregiver education.

Subarachnoid Hemorrhage

We found 10 studies focused on racial/ethnic disparities, 3 looking at sex disparities, and 4 looking at both in the diagnosis and management of SAH. Seven focused on time to treatment, and 1 focused on time to admission.

Diagnostic Assessment of SAH

An administrative claims study of more than 3,500,000 patient records from 2006 to 2012 found that among patients discharged to home from the ED with a nonspecific diagnosis of headache, male sex and non-Hispanic White race were associated with a significantly higher incidence of returning with a life-threatening neurologic condition.^{e1} SAH was the second most common missed serious neurologic condition (after ischemic stroke), while ICH was the seventh most common.

Medical Measures to Prevent Rebleeding After SAH

A small study from the Greater Cincinnati/Northern Kentucky Stroke Study found no racial, ethnic, or sex disparities in hyperacute blood pressure management after SAH in a cohort of 94 patients.^{e2} No studies addressed disparities in antifibrinolytic therapy.

Surgical and Endovascular Methods of Treatment of Ruptured Cerebral Aneurysms

Nine studies probed disparities in aneurysm treatment with 6 studies focused on racial and ethnic disparities and 3 on sex disparities. Five studies considering racial and ethnic disparities in time to SAH treatment yielded conflicting results. Two studies found longer delays to aneurysm treatment for racial and ethnic minorities compared with those for White patients,^{e3,e4} while 3 other studies did not.^{e5-e7} The 2 studies that reported disparities in time to SAH treatment both analyzed the NIS. The first examined 78,070 patients from 2002 to 2010 who underwent either clip placement or coil embolization.^{e3} Using a multivariable model, they found that Hispanic patients had a 46% increased likelihood of delayed clipping or coiling compared with White patients. Disparities were not found for other races or ethnicities. The second study assessed disparities in 7,109 patients who underwent endovascular coiling procedures from 2002 to 2011.^{e4} They found that Hispanic patients and Asian Pacific patients, but not Black patients or Native American patients, were more likely to have delays in treatment compared with White patients.

Another study from the NIS evaluated for disparities in the treatment of Hispanic patients in southwestern border states (California, Arizona, New Mexico, and Texas). In a cohort of 18,368 patients, they found that Hispanic patients in border states were more likely than non-Hispanic White patients to undergo surgical clipping rather than coiling as a treatment for SAH. This difference was not present in nonborder states.^{e8}

NIS data from 1998 to 2007 found potential racial disparities in the proportion of patients treated through clipping vs coiling. In a cohort of 20,134 patients, White patients had a higher likelihood of coil embolization than Black patients from 1998 to 2002. After the publication of the International Subarachnoid Aneurysm Trial in 2002, which found improved 1-year outcomes for those treated by endovascular coiling,^{e9} the difference decreased but did not disappear.^{e10}

Regarding sex disparities, 2 studies analyzing data from the NIS found that female patients were treated earlier than male

patients.^{e4,e11} Regarding the decision to clip vs coil, data from NIS suggest that male patients were more likely than female patients to be treated by coil embolization until 2002 when this difference dissipated.^{e10} A study analyzing 21,047 discharges from NIS 2003–2008 found that female patients were more likely than male patients to have an aneurysm clipped or coiled after SAH.^{e12} Disparities in advanced aneurysm management techniques and rates of complete aneurysm obliteration have not been evaluated.

Hospital Characteristics and Systems of Care

A study of 4,373 patients from the NIS that controlled for hospital and patient characteristics found Black patients and Hispanic patients were less likely to be transferred than White patients.^{e13} In the same analysis, there was no significant difference by sex in transfer status. The study did not account for the volume of patients with SAH treated as a hospital characteristic.

Withdrawal of Life-Sustaining Therapies, Palliative Care

We found conflicting evidence in the evaluation of differences in the withdrawal of life-sustaining therapies. A multivariate-adjusted analysis of 8,912 patients from the NIS from 2002 to 2010 found that female patients were more likely than male patients to have care withdrawn after SAH. Patients in whom care was withdrawn were less likely to be Black or Hispanic than White individuals in the same analysis.^{e14} A prospectively enrolled observational study of 1,255 patients at a single center found that Hispanic patients were less likely than White patients to receive comfort-measures-only orders, after adjusting for age and Hunt-Hess grade.^{e15} There were no significant sex differences. A different prospective observational single-center study of 1,134 patients did not find any racial, ethnic, or sex differences in the withdrawal of life-sustaining therapies.^{e16}

Rehabilitation and Recovery

Data from 31,631 admissions from the NIS from 2005 to 2010 indicated that Black patients were more likely than White patients to be discharged to postacute care despite a similar risk of death.^{e17} White, Hispanic, Asian/Pacific Islander, and Native American patients were all discharged to postacute care facilities at similar rates.

Discussion

In our scoping review of racial, ethnic, and sex disparities in the diagnosis and management of hemorrhagic stroke among adult patients in the United States, several themes emerged. First, hemorrhagic stroke disparities remain to be elucidated in key aspects of care. Second, racial and ethnic disparities may exist in long-term blood pressure management after ICH, which may contribute to increased recurrence rates among Black and Hispanic patients. Third, both racial and ethnic differences and sex differences in end-of-life care likely exist in both forms of hemorrhagic stroke. Finally, there is a paucity of

research on sex disparities in the diagnosis and management of hemorrhagic stroke, especially in SAH. Given the high incidence of SAH in women, it is important to investigate whether there are any sex disparities.

The relative lack of disparity research in hemorrhagic stroke is likely partially explained by its low incidence. Other factors may contribute. While the management of acute ischemic stroke is delineated by guidelines with strong recommendations based on high-quality evidence,^{e18} the optimal management of hemorrhagic stroke patients remains less certain.^{e19} Clear quality metrics were only just released for ICH and are yet to be released for SAH.^{e20} In the AHA/ASA guidelines for ICH, few are guided by high-level evidence. For SAH, Level A evidence guides only 2/53 recommendations. Critical process measures have not been clearly identified, and thus, disparities in process measures may be understudied. The absence of consensus surrounding the management of patients with hemorrhagic stroke may increase disparities. Weak evidence leads to uncertainty, which promotes clinician discretion and opportunities for bias.^{e21} Higher levels of evidence clarify best practices and may reduce disparity. Before the publication of the International Subarachnoid Aneurysm Trial study, White patients were more likely than Black patients to have their aneurysms coiled, but this disparity decreased after the publication of high-quality data.^{e10}

One notable area of therapeutic disparity despite strong evidence is blood pressure management to prevent recurrence of ICH. Black patients and Hispanic patients continue to have higher blood pressure at long-term follow-up after ICH compared with White patients.^{47,49,50} Persistent hypertension after ICH contributes to racial and ethnic disparities in ICH recurrence rates⁴⁷ and is partially explained by inadequate antihypertensive prescribing.⁴⁹ Racial and ethnic disparities in hypertension are not unique to patients after ICH. The causes are multifactorial and include health care access, other social determinants of health, and biologic differences.^{e22} In the case of post-ICH hypertension, presentation with such a catastrophic event should ameliorate issues of awareness and access. Innovative approaches should be pursued such as telehealth models, targeted stroke clinics, or health coaching embedded in community settings as secondary prevention of persistent hypertension and recurrent hemorrhagic stroke.^{e23-e25}

The best studied area in hemorrhagic stroke disparities or differences is the receipt or withholding of life-sustaining therapies and palliative care services. As in cohorts of patients with acute myocardial infarction or more general cohorts of hospitalized patients,^{e26} Black patients (or their surrogates) consistently opt for more aggressive treatments and less care limitations than White patients. This preference may protect Black patients from self-fulfilling prophecies whereby erroneous poor prognoses drive poor outcomes through premature withdrawal of care.³³ A Hispanic paradox, whereby Hispanic individuals “paradoxically” have similar or better

health outcomes than non-Hispanic White individuals despite lower socioeconomic status, has been previously described and attributed to self-selection in migration patterns; however, this explanation does not adequately explain the paradox.^{e27} A preference for more aggressive care in Hispanic patients and their surrogates may contribute.

The reasons for racial differences in care limitations or acceptance of palliative care remain uncertain. Authors have suggested a combination of several factors including differences in spirituality, health care system distrust, and beliefs about dying.^{e28} Much of the variation in DNR utilization, however, is not reflective of patient preference. Significant variation occurs on the hospital and regional level.^{e29} Thus, some of the differences may reflect true disparity as opposed to culturally appropriate care. More qualitative studies to evaluate differences in end-of-life care in hemorrhagic stroke would be a logical next step in understanding disparities.

Female patients may be more likely than men to receive DNR orders and palliative care after hemorrhagic stroke, although studies do exist to the contrary.²⁶ Data from other catastrophic disease states such as cardiac arrest support the supposition.^{e30} A survey of 1,648 adults did not find sex difference in the acceptance of life-prolonging treatments at end of life, suggesting that disparities in DNR orders are not driven by patient preference, but instead by medical providers or patient surrogates.^{e31} Patient comorbidities may also contribute to disparity of care, such as older age and greater frequency of cognitive impairment in female patients presenting with ICH when compared with men.

Our study has several limitations. First, many included studies came from large administrative databases, which lack data regarding important confounders. Single-center studies included such data but may not be generalizable to larger populations. Prospectively collected data from the ERICH study, which comprised 9 of the 40 reviewed ICH studies, bridge this gap but can be biased from self-selection into a study whose purpose is to study racial and ethnic variation and a Hawthorne effect. A corollary to ERICH is not available for SAH. Second, we could not adhere to best practices in disparities research because many studies were published before disparities research guidelines were available.¹² We could not always describe how race and ethnicity were categorized because those details were not available. Some studies conflated race and ethnicity. Patients could not be classified into multiple categories. Important racial and ethnic groups, such as American Indians and Alaska Natives, were ignored or subsumed into larger population groups. Substantial heterogeneity exists within racial and ethnic groups that are treated as a single category but span vast differences in geography, language, ancestry, and lived experience such as Hispanic/Latino or Asian. Key contributions from Hawaii illuminate this fallacy because they found disparities in different racial and ethnic groups that would be undiscovered had they been lumped together as “Asian.”^{e32} Third, descriptors of structural racism were largely absent despite their key role in perpetuating

health inequities.^{e30} Fourth, sex and gender were conflated, and we assumed that studies reporting gender differences were actually reporting sex differences. Conflation of sex and gender favors biologic explanations based on sex traits that may be less relevant to health care disparities than systemic discrimination based on the societal roles occupied by patients that identify as women. We found no studies explicitly evaluating gender identity disparities. By equating sex and gender and presuming a gender binary, the available literature fails to address patients with hemorrhagic stroke that are not cisgender or that are nonbinary, limiting its generalizability.

Disparities in the care of hemorrhagic stroke patients exist, especially in long-term blood pressure management and end-of-life care. More work is necessary to evaluate other aspects of care, elucidate causes, and develop solutions.

Study Funding

No targeted funding reported.

Disclosure

L. Kalasapudi, S. Williamson, A.G. Shipper, M. Motta, C. Esenwa, and F.O. Otite report no disclosures relevant to the manuscript. S. Chaturvedi serves as a consultant for Astra Zeneca, an Associate Editor for *Stroke*, and on the Editorial Board for *Neurology*[®] and *Journal of Stroke and Cerebrovascular Disease*. N.A. Morris serves on the Editorial Board for *Neurology*. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication History

Received by *Neurology* September 13, 2022. Accepted in final form March 28, 2023. Submitted and externally peer reviewed. The handling editor was Editor-in-Chief José Merino, MD, MPhil, FAAN.

Appendix Authors

Name	Location	Contribution
Lakshman Kalasapudi, BA, MA	Department of Neurology, University of Maryland School of Medicine, Baltimore	Drafting/revision of the article for content, including medical writing for content; major role in the acquisition of data; study concept or design; and analysis or interpretation of data
Stacey Williamson, MD	Department of Neurology, Henry Ford Health System, Detroit, MI	Drafting/revision of the article for content, including medical writing for content; major role in the acquisition of data
Andrea G. Shipper, MSLIS	Health Sciences and Human Services Library, University of Maryland School of Medicine, Baltimore	Major role in the acquisition of data; study concept or design
Melissa Motta, MD	Department of Neurology, University of Maryland School of Medicine; Program in Trauma, Shock Trauma Hospital, Baltimore, MD	Drafting/revision of the article for content, including medical writing for content; major role in the acquisition of data; and analysis or interpretation of data

Appendix (continued)

Name	Location	Contribution
Charles Esenwa, MD	Department of Neurology, Montefiore Medical Center, Albert Einstein College of Medicine, Bronx, NY	Drafting/revision of the article for content, including medical writing for content; analysis or interpretation of data
Fadar Oliver Otite, MD	Department of Neurology, State University of New York Upstate Medical University, Syracuse	Drafting/revision of the article for content, including medical writing for content; analysis or interpretation of data
Seemant Chaturvedi, MD	Department of Neurology, University of Maryland School of Medicine, Baltimore	Drafting/revision of the article for content, including medical writing for content; analysis or interpretation of data
Nicholas A. Morris, MD	Department of Neurology, University of Maryland School of Medicine; Program in Trauma, Shock Trauma Hospital, Baltimore, MD	Drafting/revision of the article for content, including medical writing for content; major role in the acquisition of data; study concept or design; and analysis or interpretation of data

References

- Cruz-Flores S, Rabinstein A, Biller J, et al. Racial-ethnic disparities in stroke care: the American experience: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2011;42(7):2091-2116.
- Bushnell C, Howard VJ, Lisabeth L, et al. Sex differences in the evaluation and treatment of acute ischaemic stroke. *Lancet Neurol*. 2018;17(7):641-650.
- Ovbiagele B. HEADS-UP: understanding and problem-solving: seeking hands-down solutions to major inequities in stroke. *Stroke*. 2020;51(11):3375-3381.
- Virani SS, Alonso A, Benjamin EJ, et al. Heart Disease and Stroke Statistics: 2020 update: a report from the American Heart Association. *Circulation*. 2020;141(9):e139-e596.
- Waziry R, Heshmatollah A, Bos D, et al. Time trends in survival following first hemorrhagic or ischemic stroke between 1991 and 2015 in the Rotterdam study. *Stroke*. 2020;51(3):824-829.
- Xian Y, Holloway RG, Smith EE, et al. Racial/ethnic differences in process of care and outcomes among patients hospitalized with intracerebral hemorrhage. *Stroke*. 2014;45(11):3243-3250.
- Bako AT, Pan A, Potter T, et al. Contemporary trends in the nationwide incidence of primary intracerebral hemorrhage. *Stroke*. 2022;53(3):e70-e74.
- Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol*. 2018;18(1):143.
- Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med*. 2018;169(7):467-473.
- Greenberg SM, Ziai WC, Cordonnier C, et al. 2022 Guideline for the management of patients with spontaneous intracerebral hemorrhage: a guideline from the American Heart Association/American Stroke Association. *Stroke*. 2022;53(7):e282-e361.
- Connolly ES Jr, Rabinstein AA, Carhuapoma JR, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2012;43(6):1711-1737.
- Breathett K, Spatz ES, Kramer DB, et al. The groundwater of racial and ethnic disparities research: a statement from circulation: cardiovascular quality and outcomes. *Circ Cardiovasc Qual Outcomes*. 2021;14(2):e007868.
- Karve SJ, Balkrishnan R, Mohammad YM, Levine DA. Racial/ethnic disparities in emergency department waiting time for stroke patients in the United States. *J Stroke Cerebrovasc Dis*. 2011;20(1):30-40.
- Su CM, Warren A, Kraus C, et al. Lack of racial and ethnic-based differences in acute care delivery in intracerebral hemorrhage. *Int J Emerg Med*. 2021;14(1):6.
- Tormoehlen LM, Blatsioris AD, Moser EA, et al. Disparities and guideline adherence in drugs of abuse screening in intracerebral hemorrhage. *Neurology*. 2017;88(3):252-258.
- Osman S, Zhu Z, Farag M, et al. Intracerebral hemorrhage: who gets tested for methamphetamine use and why might it matter? *BMC Neurol*. 2020;20(1):392.
- Doelfel SR, Kalagara R, Han EJ, et al. Gender disparities in stroke code activation in patients with intracerebral hemorrhage. *J Stroke Cerebrovasc Dis*. 2021;30(12):106119.
- Adeoye O, Haverbusch M, Woo D, et al. Is ED disposition associated with intracerebral hemorrhage mortality? *Am J Emerg Med*. 2011;29(4):391-395.

19. Koch S, Elkind MS, Testai FD, et al. Racial-ethnic disparities in acute blood pressure after intracerebral hemorrhage. *Neurology*. 2016;87(8):786-791.
20. Guerrero WR, Gonzales NR, Sekar P, et al. Variability in the use of platelet transfusion in patients with intracerebral hemorrhage: observations from the ethnic/racial variations of intracerebral hemorrhage study. *J Stroke Cerebrovasc Dis*. 2017;26(9):1974-1980.
21. Baharoglu MI, Cordonnier C, Al-Shahi Salman R, et al. Platelet transfusion versus standard care after acute stroke due to spontaneous cerebral haemorrhage associated with antiplatelet therapy (PATCH): a randomised, open-label, phase 3 trial. *Lancet*. 2016;387(10038):2605-2613.
22. Garcia RM, Prabhakaran S, Richards CT, Naidech AM, Maas MB. Race, socioeconomic status, and gastrostomy after spontaneous intracerebral hemorrhage. *J Stroke Cerebrovasc Dis*. 2020;29(2):104567.
23. Faigle R, Bahouth MN, Urrutia VC, Gottesman RF. Racial and socioeconomic disparities in gastrostomy tube placement after intracerebral hemorrhage in the United States. *Stroke*. 2016;47(4):964-970.
24. Cruz-Flores S, Rodriguez GJ, Chaudhry MRA, et al. Racial/ethnic disparities in hospital utilization in intracerebral hemorrhage. *Int J Stroke*. 2019;14(7):686-695.
25. Hwang DY, George BP, Kelly AG, Schneider EB, Sheth KN, Holloway RG. Variability in gastrostomy tube placement for intracerebral hemorrhage patients at US hospitals. *J Stroke Cerebrovasc Dis*. 2018;27(4):978-987.
26. Guha R, Boehme A, Demel SL, et al. Aggressiveness of care following intracerebral hemorrhage in women and men. *Neurology*. 2017;89(4):349-354.
27. Faigle R, Carrese JA, Cooper LA, Urrutia VC, Gottesman RF. Minority race and male sex as risk factors for non-beneficial gastrostomy tube placements after stroke. *PLoS One*. 2018;13(1):e0191293.
28. Sheth KN, Martini SR, Moomaw CJ, et al. Prophylactic antiepileptic drug use and outcome in the ethnic/racial variations of intracerebral hemorrhage study. *Stroke*. 2015;46(12):3532-3535.
29. Naidech AM, Toledo P, Prabhakaran S, Holl JL. Disparities in the use of seizure medications after intracerebral hemorrhage. *Stroke*. 2017;48(3):802-804.
30. Faigle R, Marsh EB, Llinas RH, Urrutia VC, Gottesman RF. Race-specific predictors of mortality in intracerebral hemorrhage: differential impacts of intraventricular hemorrhage and age among Blacks and Whites. *J Am Heart Assoc*. 2016;5(8):e003540.
31. Chen CJ, Ding D, Ironside N, et al. Predictors of surgical intervention in patients with spontaneous intracerebral hemorrhage. *World Neurosurg*. 2019;123:e700-e708.
32. Ormseth CH, Falcone GJ, Jasak SD, et al. Minority patients are less likely to undergo withdrawal of care after spontaneous intracerebral hemorrhage. *Neurocrit Care*. 2018;29(3):419-425.
33. Hemphill JC, 3rd, Newman J, Zhao S, Johnston SC. Hospital usage of early do-not-resuscitate orders and outcome after intracerebral hemorrhage. *Stroke*. 2004;35(5):1130-1134.
34. Fraser SM, Torres GL, Cai C, Choi HA, Sharief A, Chang TR. Race is a predictor of withdrawal of life support in patients with intracerebral hemorrhage. *J Stroke Cerebrovasc Dis*. 2018;27(11):3108-3114.
35. Murthy SB, Moradiya Y, Hanley DF, Ziai WC. Palliative care utilization in non-traumatic intracerebral hemorrhage in the United States. *Crit Care Med*. 2016;44(3):575-582.
36. Lillemoe K, Lord A, Torres J, Ishida K, Zeisler B, Lewis A. Factors associated with DNR status after nontraumatic intracranial hemorrhage. *Neurohospitalist*. 2020;10(3):168-175.
37. Zahuranec DB, Brown DL, Lisabeth LD, et al. Ethnic differences in do-not-resuscitate orders after intracerebral hemorrhage. *Crit Care Med*. 2009;37(10):2807-2811.
38. Zurasky JA, Aiyagari V, Zazulia AR, Shackelford A, Diringer MN. Early mortality following spontaneous intracerebral hemorrhage. *Neurology*. 2005;64(4):725-727.
39. Faigle R, Ziai WC, Urrutia VC, Cooper LA, Gottesman RF. Racial differences in palliative care use after stroke in majority-white, minority-serving, and racially integrated U.S. hospitals. *Crit Care Med*. 2017;45(12):2046-2054.
40. McFarlin J, Hailey CE, Qi W, et al. Associations between patient characteristics and a new, early do-not-attempt resuscitation order after intracerebral hemorrhage. *J Palliat Med*. 2018;21(8):1161-1165.
41. Patel AA, Mahajan A, Benjo A, et al. A national perspective of do-not-resuscitate order utilization predictors in intracerebral hemorrhage. *Neurohospitalist*. 2016;6(1):7-10.
42. Suolond D, Chen BJ, Faigle R. Temporal trends in racial and ethnic disparities in palliative care use after intracerebral hemorrhage in the United States. *Stroke*. 2022;53(3):e85-e87.
43. Shaw KM, Gallek MJ, Sheppard KG, et al. Ethnic differences in withdrawal of life support after intracerebral hemorrhage. *Hawaii J Med Public Health*. 2015;74(6):203-209.
44. Ganti L, Jain A, Yerragondou N, et al. Female gender remains an independent risk factor for poor outcome after acute nontraumatic intracerebral hemorrhage. *Neurol Res Int*. 2013;2013:219097.
45. Nakagawa K, Vento MA, Seto TB, et al. Sex differences in the use of early do-not-resuscitate orders after intracerebral hemorrhage. *Stroke*. 2013;44(11):3229-3231.
46. Craen A, Mangal R, Stead TG, Ganti L. Gender differences in outcomes after non-traumatic intracerebral hemorrhage. *Cureus*. 2019;11(10):e5818.
47. Rodriguez-Torres A, Murphy M, Kourkoulis C, et al. Hypertension and intracerebral hemorrhage recurrence among white, black, and Hispanic individuals. *Neurology*. 2018;91(1):e37-e44.
48. Leasure A, King Z, Torres-Lopez V, et al. Racial/ethnic disparities in the risk of intracerebral hemorrhage recurrence. *Neurology*. 2019;92(15):e314-e322.
49. Biffi A, Teo KC, Castello JP, et al. Impact of uncontrolled hypertension at 3 months after intracerebral hemorrhage. *J Am Heart Assoc*. 2021;10(11):e020392.
50. Zahuranec DB, Wing JJ, Edwards DF, et al. Poor long-term blood pressure control after intracerebral hemorrhage in a predominantly african american hypertensive population. *Stroke*. 2012;43(2):2580-2585.

eReferences are listed at [links.lww.com/WNL/C825](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC825).